

# サクロン (顆粒)

Category-2 OTC Drug



## サクロン®

### Saclon Granule

A green stomach medicine that refreshes the stomach. For stomach retching, heartburn, and excessive drinking.

Excessive secretion of gastric acid may cause stomach retching and heartburn.

As another cause, alcohol may directly irritate the stomach mucous membrane itself due to excessive drinking.

Saclon attenuates (neutralizes) the action of excess gastric acid, suppresses its secretion, and restores and protects the damaged stomach mucous membrane with a green ingredient made from chlorophyll.

For people who are suffering stomach retching, gripping pain, and nausea when brushing their teeth in the morning. Take Saclon, which will give you refreshing effects.



## Indications



Stomach  
heaviness



heartburn



stomachache



hangover

Heartburn, excessive drinking, stomachache, hyperacidity, stomach heaviness, feeling of discomfort in the stomach, feeling of fullness in the stomach, heaviness in the stomach, heaviness in the chest, belching, nausea (retching, stomach retching, retching due to hangovers and overdrinking, sick feeling, and urge to vomit), vomiting



# Dosage and Administration

Take the following dose with cold or hot water between meals and before going to bed on an empty stomach.

Age	Dosage	Number of doses
Adults (15 years or over)	1 sachet	3 times per day
8 years or over and under 15 years	1/2 sachets	3 times per day
Under 8 years	Do not take this medicine	-

\* "Between meals" means the time where you have an empty stomach with almost no food about 2 hours after meals.

## Precautions for Dosage and Administration

When allowing children (aged 8 years or over and under 15 years) to take the medicine, a guardian must be present to watch the child and instruct them how to take it.



# Ingredients

## Ingredients and their actions

The daily adult dose of 3 sachets (3.84 g) contains the following ingredients.

Ingredients	Amount	Action
Potassium copper chlorophyllin	120 mg	It restores and protects the damaged stomach mucous membrane.
Anhydrous dibasic calcium phosphate	1,020 mg	It directly neutralizes the excess gastric acid.
Precipitated calcium carbonate	1,020 mg	It directly neutralizes the excess gastric acid.
Magnesium hydroxide	960 mg	It directly neutralizes the excess gastric acid.
Scopolia extract	30 mg	It suppresses the secretion of gastric acid and stops the pain.

## Additives

---

Corn starch, hydroxypropyl cellulose, l-menthol, flavor, potassium chloride, cinnamon bark, magnesium stearate, povidone

## Precautions for Ingredients

---

1. The stool may turn green after taking this drug. This is due to excretion of the ingredient potassium copper chlorophyllin (green). It is not a cause for concern.
2. Saclon does not use aluminum-containing ingredients.



## Precautions

### ⚠ Precautions for Use

---

#### When not to use the product

[If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.]

1. This drug should not be taken together with the following drugs:

Gastrointestinal pain relief/antispasmodic drugs

2. Nursing women should either not take this medicine or stop nursing while taking this medicine.  
(This drug is excreted in breast milk and may cause a fast pulse in nursing infants.)

---

## Consultation

**1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before taking this drug.**

1. Patients undergoing medical treatment from a physician
2. Pregnant women or women who believe they may be pregnant
3. The elderly
4. Patients who have experienced allergic symptoms associated with drugs etc.
5. Persons with the following symptom:  
Difficulty urinating
6. Persons diagnosed as having the following:  
Kidney disease, heart disease, glaucoma, thyroid disease

**2. If the following symptoms are observed after taking this drug, these may be adverse reactions, so immediately discontinue the use of this drug and show this instruction leaflet to your physician, pharmacist, or registered salesperson for a consultation.**

Parts Affected	Symptom
Skin	Rash/redness, itching

**3. The following symptoms may be observed after taking this drug. If these symptoms persist or worsen, discontinue the use of this drug and show this instruction leaflet to your physician, pharmacist, or registered salesperson for a consultation.**

Dry mouth, constipation, diarrhea

**4. If symptoms do not improve even after taking this medicine for about 2 weeks, stop taking this medicine and consult a physician, pharmacist, or registered salesperson, being sure to take this instruction leaflet with you.**

---

## Other precautions

Some may have trouble lactating.

---

## Precautions for storage and handling

**1. Store in a cool, low humidity place away from direct sunlight.**

2. Keep out of reach of children.
3. Do not transfer to a different container. Moreover, do not put other drugs in this container. (This may lead to misuse or alter the quality of the drug.)
4. When taking the granules remaining in a sachet divided in multiple portions, fold back the opening of the sachet before storing it and do not take the granules stored for more than 2 days.
5. Do not use the product after the expiration date.



## Packaging and Price

Packaging	Recommended retail price
10 sachets	1,078yen, tax included (980yen, tax excluded)
20 sachets	2,046yen, tax included (1,860yen, tax excluded)
32 sachets	3,113yen, tax included (2,830yen, tax excluded)