

# チョコラBBゴールドリッチ （製造終了）

Designated quasi-drug



## チョコラBB<sup>®</sup>ゴールドリッチ

Quick acting care for tiredness with general malaise and poor skin condition!

- Contains 16 ingredients, the most in the Chocora BB series. In addition to B group vitamins and 1500 mg of taurine, it contains a luxurious blend of 6 crude drugs that also appeal to women.
- Has a flavor of bergamot, which is also used in Earl Grey tea and aroma oils. We have chosen a flavor that is familiar to women and pleasant to drink.

## Indications



fatigue and  
malaise



loss of  
appetite

- Maintaining and improving physical strength, physical resistance, or mental concentration

- Recovering from and preventing fatigue
- Weak constitution (including physical weakness due to aging)Improving or preventing poor physical condition due to
- Improving or preventing poor physical condition due to poor nutrition in daily life: becoming tired easily, residual tiredness, lack of physical strength, heaviness in the body, physical malaise, poor skin condition (skin roughness or dryness), sensitivity to cold - poor blood circulation
- Nutritional supplementation when physical strength is impaired during or after illness, during exhaustion due to fever, for loss of appetite, during pregnancy or lactation (or before and after labor), etc.



## Dosage and Administration

For adults (15 years and older), take 1 bottle (50 mL) once a day.

Follow the dosage and administration method. (If using this medicine at the same time as other products with ingredients such as vitamins, be careful not to take excessive amounts)



## Active Ingredients (in 1 bottle of 50 mL)


Ingredients	Amount
Vitamin B <sub>2</sub> phosphate ester	15 mg
Vitamin B <sub>6</sub>	10 mg
Vitamin B <sub>1</sub> nitrate	10 mg
Royal jelly tincture	300 mg (300 mg as royal jelly)
Taurine	1500 mg
Glycine	50 mg
L-Arginine hydrochloride	150 mg
L-Lysine hydrochloride	100 mg

Ginseng extract	54 mg(600 mg as ginseng)
Touki (Japanese angelica root) fluidextract	0.2 mL (200 mg as Touki)
Shokyo (ginger) extract	38.8 mg (400 mg as Shokyo)
Ryuganniku (longan aril) extract	90.9 mg (300 mg as Ryuganniku)
Kukoshi (lycium fruit) fluidextract	0.2 mL (200 mg as Kukoshi)
Nicotinamide	40 mg
d- $\alpha$ -Tocopherol acetate (natural vitamin E)	10 mg
Anhydrous caffeine	50 mg

## Inactive Ingredients

Sodium benzoate, citric acid, sodium citrate, glycerin, ethyl p-hydroxybenzoate, polyoxyethylene hydrogenated castor oil, flavor, acesulfame-K, sucrose esters of fatty acids, sucralose, polyglyceryl monostearate, medium chain fatty acid triglyceride, dextrin, trehalose, white soft sugar, povidone, starch syrup, DL-malic acid (alcohol content no more than 0.49 mL)

Taking this medicine may cause urine to turn yellow, but this is due to assimilation of the vitamin B<sub>2</sub> contained in the medicine, with part of it being excreted in the urine. It is not a cause for concern.



## Precautions

### Precautions for Use



### Consultation

1. If the following symptoms are observed after taking this medicine, these may be adverse reactions, so immediately discontinue the use of this medicine, and consult a physician, pharmacist, or registered salesperson, being sure to take this medicine with you.

Parts Affected	Symptom
Skin	Rash
Digestive organs	Stomach discomfort

- The following symptoms may be observed after taking this medicine. If these symptoms persist or worsen, discontinue the use of this medicine, and consult a physician, pharmacist, or registered salesperson, being sure to take this medicine with you.

Diarrhea

- When symptoms do not improve even after using the medicine for a while, discontinue the use of this medicine, and consult a physician, pharmacist, or registered salesperson, being sure to take this medicine with you.

## Precautions for storage and handling

- Store sealed in a cool place away from direct sunlight.
- Keep out of reach of children.
- Do not transfer to a different container. (It may lead to misuse or alter the quality.)
- Due to the crude drug extracts contained in this medicine, cloudiness may be observed; however, there is no change in the effects.
- Do not take medicine that has exceeded its expiration date.



## Packaging and Price

Packaging	Recommended retail price
50 mL	548yen, tax included (498yen, tax excluded)
50 mL × 3	1,643yen, tax included (1,494yen, tax excluded)
50 mL × 10	5,478yen, tax included (4,980yen, tax excluded)