

本製品は日本国法に基づき製造販売されたものです。  
 製品および文書は日本国外の法規に準じているわけではありません。  
 本文書は、日本語の製品情報を翻訳した文書です。使用前に必ずお読みください。  
 また、必要な時に読めるように大切に保管してください。  
 This product has been manufactured and sold based on Japanese law.  
 Neither the product nor this informational leaflet will necessarily conform to the laws of countries outside of Japan.  
 This leaflet contains a translation of product information from Japanese. You should read this written explanation before using the product.  
 Store this informational leaflet safely so that it can be read when necessary.



Discovering everything from prevention to cure  
 Eisai.jp

Category-3 OTC Drug

For eczema and stomatitis in children (3 months and older).

## チョコラBB®こどもシロップ

### Chocola BB Children's Syrup

Vitamin B<sub>2</sub> has been called the “vitamin for fat metabolism” and the “vitamin for skin and mucosa.” Deficiencies of this vitamin can cause symptoms such as fatigue, eczema, dermatitis and stomatitis to occur easily.

チョコラBB こどもシロップ contains vitamin B<sub>2</sub> as its main ingredient, along with vitamins B<sub>1</sub> and B<sub>6</sub> and biotin, which is known as vitamin for anti-dermatitis. It is a syrup form for children to swallow easily.



Package	Recommended retail price (tax included)
30 mL × 2	¥756

Related website



[Chocola.com](http://Chocola.com) (in Japanese)

### Indications

**Relief of the following symptoms:**

Skin roughness, rashes, sores, acne, eczema, dermatitis, stomatitis, angular stomatitis, canker sores, glossitis, rosacea, congestion of the eyes, itchy eyes

Please consult with a physician or pharmacist if no improvement is seen in these symptoms after a month of use.

**Supplementation of vitamin B<sub>2</sub> in the following cases:**

When physically fatigued, or when strength is decreased during and after illness

### Dosage and Administration

Take twice a day after meals, using the measuring cup provided to measure the dose

Age	Dosage	Number of doses
Between 7 and 15 years	15 mL	Twice a day
Between 1 and 7 years	7.5 mL	
Between 3 months and 1 year	5 mL	
Under 3 months	Do not take this medicine	

**Precautions for Dosage and Administration**

When allowing children (3 months–15 years) to take the medicine, a guardian must be present to watch and tell them how to take it.

## Active Ingredients and Actions

One 30 mL bottle contains the following active ingredients.

Ingredient	Amount	Action
Riboflavin Sodium Phosphate (Vitamin B <sub>2</sub> phosphate ester)	19 mg	<ul style="list-style-type: none"><li>Supports healthy skin and mucosa.</li><li>Aids fat metabolism, converting it to energy.</li></ul>
Pyridoxine Hydrochloride (Vitamin B <sub>6</sub> )	12 mg	<ul style="list-style-type: none"><li>Supports healthy skin and mucosa.</li><li>Aids protein metabolism, converting it to energy.</li></ul>
Thiamine Nitrate (Vitamin B <sub>1</sub> nitrate)	6 mg	<ul style="list-style-type: none"><li>Aids carbohydrate metabolism, converting it to energy.</li></ul>
Nicotinamide	20 mg	<ul style="list-style-type: none"><li>Supports healthy skin and mucosa.</li></ul>
Biotin	250 µg	<ul style="list-style-type: none"><li>Supports healthy skin and mucosa and suppresses inflammation of the skin.</li></ul>

### Inactive Ingredients

Sodium benzoate, ethanol (0.1 mL or less), citric acid, sodium citrate, D-sorbitol, paraben, propylene glycol, flavor, hydrogenated maltose starch syrup, glycine, propyl gallate, povidone, DL-malic acid, pH adjuster

### Precautions for Ingredients

Taking this medicine may cause urine to turn bright yellow, but this is due to assimilation of the vitamin B2 contained in the medicine, with part of it being excreted in the urine. It is not a cause for concern.



## Precautions

### Consultation

- If the following symptoms are observed after taking this medicine, these may be adverse reactions. Immediately discontinue the use of this medicine and show this informational leaflet to a physician, pharmacist or registered salesperson for a consultation.

Parts Affected	Symptom
Skin	Rash
Digestive organs	Stomach discomfort

- The following symptom may be observed after taking this medicine. If this symptom persists or worsens, discontinue the use of this medicine and show this informational leaflet to a physician, pharmacist or registered salesperson for a consultation.

Diarrhea

- If symptoms do not improve even after taking the medicine for about 1 month, stop taking the medicine, and consult a physician, pharmacist or registered salesperson, bringing this informational leaflet with you.

## Precautions for Storage and Handling

- Store sealed in a cool place away from direct sunlight.
- Keep out of reach of children.
- Keep refrigerated after opening the bottle, and use it as soon as possible.
- Do not transfer to a different container. Also, do not transfer other medications, etc., into this container. (It may lead to misuse or alter the quality.)
- After administration, rinse the measuring cup, and store it securely with the syrup.
- Do not take medicine that has exceeded its expiration date.

Contact us: For inquiries, please complete the Contact Form. We will endeavor to respond to you as quickly as possible, although there may be some delay depending on the nature of the inquiry and other factors.

Contact Form (English only): [https://inquiry.eisai.co.jp/webapp/form/17672\\_sdab\\_2/index.do](https://inquiry.eisai.co.jp/webapp/form/17672_sdab_2/index.do)