

本製品は日本国法に基づき製造販売されたものです。
製品および文書は日本国外の法規に準じているわけではありません。
本文書は、日本語の製品情報を翻訳した文書です。使用前に必ずお読みください。
また、必要な時に読めるように大切に保管してください。
This product has been manufactured and sold based on Japanese law.
Neither the product nor this informational leaflet will necessarily conform to the laws of countries outside of Japan.
This leaflet contains a translation of product information from Japanese. You should read this written explanation before using the product.
Store this informational leaflet safely so that it can be read when necessary.



Discovering everything from prevention to cure
Eisai.jp

Designated quasi-drug

Nutritional supplementation and nutrient/tonic for physical fatigue

チョコラBB[®]ゴールドリッチ

Chocola BB Gold Rich

Quick acting care for tiredness with general malaise!
Contains 16 ingredients, the most in the Chocola BB series. In addition to B group vitamins and 1500 mg of taurine, it contains a luxurious blend of 6 crude drugs that also appeal to women. Has a flavor of bergamot, which is also used in Earl Grey tea and aroma oils. We have chosen a flavor that is familiar to women and pleasant to drink.



Package	Recommended retail price (tax included)
50 mL	¥754
50 mL × 10 bottles	¥7,538
50 mL × 50 bottles	¥37,692

Related website



Chocola.com (in Japanese)

Indications

- Nutritional supplementation for physical fatigue, during and after illness, for loss of appetite, nutritional disorders, exhaustion due to fever, during pregnancy or lactation, etc.
- For nutrients/as a tonic
- For a weak constitution

Dosage and Administration

For adults (15 years and older), take 1 bottle (50 mL) once a day.

Precautions for Dosage and Administration

Follow the dosage and administration method. (If using this medicine at the same time as other products with ingredients such as vitamins, be careful not to take excessive amounts)

Active Ingredients

Active Ingredients (in 1 bottle of 50 mL)

Ingredient	Amount
Vitamin B ₂ phosphate ester	15 mg
Vitamin B ₆	10 mg
Vitamin B ₁ nitrate	10 mg
Royal jelly tincture	300 mg (300 mg as royal jelly)
Taurine	1500 mg
Glycine	50 mg
L-Arginine hydrochloride	150 mg
L-Lysine hydrochloride	100 mg
Ginseng extract	54 mg (600 mg as ginseng)
Touki (Japanese angelica root) fluidextract	0.2 mL (200 mg as Touki)
Shokyo (ginger) extract	38.8 mg (400 mg as Shokyo)
Ryuganniku (longan aril) extract	90.9 mg (300 mg as Ryuganniku)
Kukoshi (lycium fruit) fluidextract	0.2 mL (200 mg as Kukoshi)
Nicotinamide	40 mg
d- α -Tocopherol acetate (natural vitamin E)	10 mg
Anhydrous caffeine	50 mg

Inactive Ingredients

Sodium benzoate, citric acid, sodium citrate, glycerin, ethyl p-hydroxybenzoate, polyoxyethylene hydrogenated castor oil, flavor, acesulfame-K, sucrose esters of fatty acids, sucralose, polyglyceryl monostearate, medium chain fatty acid triglyceride, dextrin, trehalose, white soft sugar, povidone, starch syrup, DL-malic acid (alcohol content no more than 0.49 mL)

Precautions for Ingredients

Taking this medicine may cause urine to turn yellow, but this is due to assimilation of the vitamin B₂ contained in the medicine, with part of it being excreted in the urine. It is not a cause for concern.



Precautions

Consultation Needed for These Cases

1. In the following cases, immediately discontinue the use of this medicine and consult a physician or pharmacist, being sure to take this medicine with you.

(1) If the following symptoms are observed after taking this medicine

Parts affected	Symptom
Skin	Rash
Digestive organs	Stomach discomfort

(2) When symptoms do not improve even after using the medicine for a while

2. The following symptoms may be observed after taking this medicine. If these symptoms persist or worsen, discontinue the use of this medicine, and consult a physician or pharmacist.

Diarrhea

Precautions for Storage and Handling

1. Store sealed in a cool place away from direct sunlight.
2. Keep out of reach of children.
3. Do not transfer to a different container. (It may lead to misuse or alter the quality.)
4. Due to the crude drug extracts contained in this medicine, cloudiness may be observed; however, there is no change in the effects.
5. Do not take medicine that has exceeded its expiration date.

Contact us: For inquiries, please complete the Contact Form. We will endeavor to respond to you as quickly as possible, although there may be some delay depending on the nature of the inquiry and other factors.

Contact Form (English only): https://inquiry.eisai.co.jp/webapp/form/17672_sdab_2/index.do