

本製品は日本国法に基づき製造販売されたものです。
製品および文書は日本国外の法規に準じているわけではありません。
本文書は、日本語の製品情報を翻訳した文書です。使用前に必ずお読みください。
また、必要な時に読めるように大切に保管してください。
This product has been manufactured and sold based on Japanese law.
Neither the product nor this informational leaflet will necessarily conform to the laws of countries outside of Japan.
This leaflet contains a translation of product information from Japanese. You should read this written explanation before using the product.
Store this informational leaflet safely so that it can be read when necessary.



Discovering everything from prevention to cure
Eisai.jp

Designated quasi-drug

Improving or preventing poor physical condition due to poor nutrition in daily life
Poor skin condition (skin roughness or dryness).

チョコラBB[®]ハイパー

Chocola BB Hyper

- For when you want to do your best!
- Contains 13 ingredients, including B group vitamins, amino acids, taurine, and also royal jelly, ginseng, and guarana.
- It has a full bodied yet refreshing taste that is pleasant for women to drink.



Package	Recommended retail price (tax included)
50 mL	¥410
50 mL × 3	¥1,231
50 mL × 10	¥4,104

Related website



Chocola.com (in Japanese)

Indications

- Maintaining and improving physical strength, physical resistance, or mental concentration
- Recovering from and preventing fatigue
- Improving or preventing poor physical condition due to a weak constitution (including weak constitutions due to aging)
- Improving or preventing poor physical condition due to poor nutrition in daily life
Becoming tired easily, residual tiredness, lack of physical strength, heaviness in the body, physical malaise
Poor skin condition (skin roughness or dryness)
Becoming cold easily, poor blood flow
- Nutritional supplementation when physical strength is impaired during or after illness, during exhaustion due to fever, for loss of appetite, and during pregnancy or lactation (or before and after labor), etc.

Dosage and Administration

Age	Dosage	Number of doses
Adult (15 years or older)	1 bottle (50 mL)	Once a day
Child (Under 15 years)	Do not take this medicine	

Precautions for Dosage and Administration

Follow the dosage and administration method. (If using this medicine at the same time as other products with ingredients such as vitamins, be careful not to take excessive amounts)

Active Ingredients

Active Ingredients (in 1 bottle of 50 mL)

Ingredient	Amount
Vitamin B ₂ phosphate ester	15 mg
Vitamin B ₆	10 mg
Vitamin B ₁ nitrate	10 mg
Royal jelly tincture (300 mg as royal jelly)	300 mg
Taurine	1200 mg
Glycine	50 mg
L-Arginine hydrochloride	150 mg
L-Lysine hydrochloride	100 mg
Guarana fluidextract (150 mg as guarana)	0.15 mL
Ginseng dry extract (600 mg as ginseng)	41.4 mg
Joteishi extract (250 mg as Joteishi)	62.5 mg
Nicotinamide	40 mg
Caffeine hydrate	45 mg

Inactive Ingredients

Sodium benzoate, ethyl vanillin, citric acid, sodium citrate, glycerin, vanillin, paraben, propylene glycol, flavor, acesulfame-K, erythritol, sucralose, refined stevia extract, dextrin, honey, propyl gallate, DL-malic acid (alcohol content no more than 0.4 mL)

Precautions for Ingredients

Taking this medicine may cause urine to turn yellow, but this is due to assimilation of the vitamin B2 contained in the medicine, with part of it being excreted in the urine. It is not a cause for concern.



Precautions

Consultation Needed for These Cases

1. In the following cases, immediately discontinue the use of this medicine and consult a physician or pharmacist, being sure to take this medicine with you.

(1) If the following symptoms are observed after taking this medicine

Parts affected	Symptom
Skin	Rash
Digestive organs	Stomach discomfort

(2) When symptoms do not improve even after using the medicine for a while

2. The following symptoms may be observed after taking this medicine. If these symptoms persist or worsen, discontinue the use of this medicine, and consult a physician or pharmacist.

Diarrhea

Precautions for Storage and Handling

1. Store sealed in a cool place away from direct sunlight.
2. Keep out of reach of children.
3. Do not transfer to a different container. (It may lead to misuse or alter the quality.)
4. Due to the crude drug extracts contained in this medicine, cloudiness may be observed; however, there is no change in the effects.
5. Do not take medicine that has exceeded their expiration date.

Contact us: For inquiries, please complete the Contact Form. We will endeavor to respond to you as quickly as possible, although there may be some delay depending on the nature of the inquiry and other factors.

Contact Form (English only): https://inquiry.eisai.co.jp/webapp/form/17672_sdab_2/index.do