

本製品は日本国法に基づき製造販売されたものです。
 製品および文書は日本国外の法規に準じているわけではありません。
 本文書は、日本語の製品情報を翻訳した文書です。使用前に必ずお読みください。
 また、必要な時に読めるように大切に保管してください。
 This product has been manufactured and sold based on Japanese law.
 Neither the product nor this informational leaflet will necessarily conform to the laws of countries outside of Japan.
 This leaflet contains a translation of product information from Japanese. You should read this written explanation before using the product.
 Store this informational leaflet safely so that it can be read when necessary.



Discovering everything from prevention to cure
 Eisai.jp

Category-3 OTC drug

Removing bad breath, hangover

サクロフィール錠

Saclophyl Tablets

サクロフィール錠 contain a green ingredient made from chlorophyll that acts on the substances in the body that cause odors, to remove bad breath.



Package	Recommended retail price (tax included)
12 tablets	¥356
50 tablets	¥1,058

Indications

Elimination of bad breath, hangover

Dosage and Administration

Take the following doses with cold or hot water.

Age	Dosage	Number of doses
Adult (15 years or older)	3 to 4 tablets	1 to 3 time(s) per day
Child (Under 15 years)	Do not take this medicine	

How to Remove the Tablets from the Packaging

For the 12 tablet package, press firmly on a raised part of the sheet containing the tablet to break the aluminum foil on the reverse side, and then take out a tablet and swallow it. (Mistakenly swallowing tablets that are still in their sheet may cause unexpected accidents such as tearing of the esophageal membrane.)

Active Ingredients and Action

One tablet contains the following active ingredients.

Ingredient	Amount	Action
Sodium copper chlorophyllin	15 mg	Acts on the substances in the body that cause odors, to remove bad breath.

Inactive Ingredients

Corn starch, dl-menthol, magnesium silicate, calcium stearate, white soft sugar

Precautions for Ingredients

1. After you take a tablet, the surface of your tongue will become green. This coloration is due to sodium copper chlorophyllin. It will disappear with time.
2. After you take a tablet, your stool may become green and have less smell than usual. This is coloration and a deodorizing action due to sodium copper chlorophyllin, and is not a cause for concern.



Precautions

Consultation

1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
Patients undergoing medical treatment from a physician
2. The following symptoms may be observed after taking this medicine. If these symptoms persist or worsen, discontinue the use of this medicine, and show this package to a physician, pharmacist, or registered salesperson for a consultation.
Soft stool

Other Precautions

If you have an illness that causes bad breath, make sure that you also receive treatment for this illness.

Precautions for Storage and Handling

1. Store (sealed) in a cool, dry place away from direct sunlight. Information in parentheses is shown on the 50 tablet package.
2. Keep out of reach of children.
3. Do not transfer to a different container. Also, do not transfer other medications, etc., into this container. (It may lead to misuse or alter the quality.)
4. Do not take tablets that have exceeded their expiration date.
5. If you are using the 50 tablet package, the following precaution applies.
The packing material inside the container is provided to prevent damage to the tablets during transport. After opening the cap, discard the packing material. Keep the container sealed during storage.

Contact us: For inquiries, please complete the Contact Form. We will endeavor to respond to you as quickly as possible, although there may be some delay depending on the nature of the inquiry and other factors.

Contact Form (English only): https://inquiry.eisai.co.jp/webapp/form/17672_sdab_2/index.do